

1 **Short title:** *Bifiguratus adelaidae*, a new member of Mucoromycotina

2 ***Bifiguratus adelaidae* gen. et sp. nov., a new member of Mucoromycotina in endophytic and soil-dwelling**
3 **habitats**

4 Terry J. Torres-Cruz

5 Terri L. Billingsley Tobias

6 Maryam Almatruk

7 *Department of Biological Sciences, Western Illinois University, Macomb, IL, USA 61455*

8 Cedar N. Hesse

9 Cheryl R. Kuske

10 *Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM, USA 87545*

11 Alessandro Desirò

12 Gian Maria Niccolò Benucci

13 Gregory Bonito

14 *Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, MI, USA 48824*

15 Jason E. Stajich

16 *Department of Plant Pathology & Microbiology, University of California, Riverside, CA, USA 92521*

17 Christopher Dunlap

18 *National Center for Agricultural Utilization Research, Agricultural Research Service, U.S. Department of*
19 *Agriculture, Peoria, IL, USA 61604*

20 A. Elizabeth Arnold

21 *School of Plant Sciences and Department of Ecology and Evolutionary Biology, The University of Arizona,*
22 *Tucson, AZ, USA 85721*

23 Andrea Porras-Alfaro¹

24 *Department of Biological Sciences, Western Illinois University, Macomb, IL, USA 61455*

¹ Corresponding author. Email: a-porras-alfaro@wiu.edu

Abstract

Illumina amplicon sequencing of soil in a temperate pine forest in the southeastern United States detected an abundant, N-responsive fungal genotype of unknown phylogenetic affiliation. Two isolates with ribosomal sequences consistent with that genotype were subsequently obtained. Examination of records in GenBank revealed that a genetically similar fungus had been isolated previously as an endophyte of moss in a pine forest in the southwestern USA. The three isolates were characterized using morphological, genomic, and multi-locus molecular data (18S, ITS, and 28S rRNA sequences). Phylogenetic and maximum likelihood phylogenomic reconstructions revealed that the taxon represents a novel lineage in Mucoromycotina, only preceded by *Calcarisporiella*, the earliest diverging lineage in the subphylum. Sequences for the novel taxon are frequently detected in environmental sequencing studies and it is currently part of UNITE's dynamic list of most wanted fungi. The fungus is dimorphic, grows best at room temperature, and is associated with a wide variety of bacteria. Here a new monotypic genus, *Bifiguratus*, is proposed, typified by *Bifiguratus adalaidae*.

KEY WORDS: Bryophyte, dimorphic, endophyte, environmental sampling, Mucoromycota, Pinaceae, soil.

INTRODUCTION

Although fungal species are being described continuously, and have been for centuries, the majority of fungi remain undiscovered. Working estimates predict that 5.1 million species of fungi exist, yet only ~100,000 species have been described formally (Blackwell 2011). A recent study on the tree of life encompassing more than 1000 uncultivated organisms highlighted the large number of organisms for which cultured representatives are unknown (Hug et al. 2016). Indeed, new sequencing techniques frequently reveal large numbers of uncharacterized taxa that

have not been described due to the lack of representative reference specimens in herbaria and/or culture collections (Martin et al. 2011). A major gap in our understanding of fungal diversity lies in determining the connections between fungi detected through next-generation sequencing or other culture-free methods, and fungi already known to mycologists (Nilsson et al. 2016; Hesse et al. 2016). One approach is to use next-generation data sets to guide subsequent efforts to isolate target fungi from the same environments, thus providing biological material for further functional and molecular analyses (Hesse et al. 2016).

Many branches of the fungal tree of life are still poorly resolved and underrepresented (or lacking) in phylogenies. This is particularly true in lineages of early diverging terrestrial fungi (including those formerly classified as Zygomycota; Spatafora et al. 2016). Improved sampling of diversity in these groups is critical for illuminating key evolutionary innovations in the fungal kingdom and assessing the ecological functioning of novel taxa or those reported previously as non-culturable (or uncultured) (Grigoriev et al. 2011).

A previous investigation of the mycobiota associated with soil in a pine forest in North Carolina, USA (Hesse et al. 2016) detected a fungal taxon by 28S nrRNA amplicon sequencing. This fungal taxon was common in the soil and responded positively to forest N fertilization treatments (Weber et al. 2013; Hesse et al. 2016). Particle filtration of soils from the same site subsequently yielded two isolates (TLT265, YP243) representing the same 28S nrRNA phylotype. When looking for similar sequences in the National Center for Biotechnology Information (NCBI), one other match (AZ0501) with >99% similarity for the ITS, 28S, and 18S regions was found. These three strains were compared and found to be similar in morphology and molecular characters. Here we report and describe a novel taxon, *Bifiguratus adelaidae*, and place it as an early diverging lineage in Mucoromycotina through multi-locus and phylogenomic

analyses. The presence of mucoid growth on the edges of the colonies suggested the presence of bacteria on the fungus, which was evaluated using culturing, microscopy, and next generation sequencing techniques.

MATERIALS AND METHODS

Sample collection and site description —

As part of a previous study, soil cores were collected from a free air CO₂ enrichment (FACE) experiment located in a *Pinus taeda* plantation in the Blackwood Division of the Duke Forest, North Carolina, USA (35°58'N, 79°06'W; elevation 130 m). The FACE site was established in 1983 (Weber et al. 2013). In 2005, the CO₂ enrichment plots were quartered and annual N-fertilization experiments began in two randomly selected quadrants in the form of ammonium nitrate pellets (11.2 g N m⁻²). Soil sampling was designed to take into account the long-term field experiment and to minimally disturb soils in the plots. The sampling was performed within the fertilized and unfertilized quadrants of ring 5 (Weber et al. 2013). In October 2013, 10 cm soil cores were collected from one N-fertilized and one unfertilized quadrant within ambient CO₂ FACE plot 5. Immediately after collection, each core was sectioned into multiple vertical strata: Oi horizon (forest floor), Oa horizon (0–2 cm deep), upper A horizon (2–5 cm deep), and lower A horizon (5–10 cm deep). Each soil fraction was homogenized in an individual zip-lock bag. Soil samples were transported back to the laboratory and stored at 4 °C until use.

Fungal isolation —

Fungi were isolated following two techniques, standard serial dilution methods for soil (Gupta et al. 2012) and a particle-pasteurization filtration method (Mueller et al. 2004). Detailed methods for isolation can be found in Hesse et al. (2016). Briefly, for the particle filtration

method, seven sieves were used to separate different particle sizes (4000, 2000, 250, 500, 125, 63, and $< 63 \mu\text{m}$). One gram of soil was loaded in the upper sieve (4000 μm). Particles were washed with 4–6 liters of autoclaved sterile water for 10 minutes. The last rinse was 1 liter of 70% ethanol to surface-sterilize the soil. The soil particles from each sieve were transferred to 9 ml of 0.2% water agar (used to suspend the particles). Dilutions of 10^{-1} and 10^{-2} were plated on peptone-yeast glucose agar supplemented with tetracycline (50 mg/L) and streptomycin (50 mg/L). Cultures were incubated at 25 °C for 15 days. Plates were checked regularly to obtain pure cultures.

Cultures and specimens from this study are preserved in the USDA-ARS Culture Collection (Peoria, Illinois), the herbarium of the University of Michigan (Ann Arbor, Michigan), the Robert L. Gilbertson Mycological Herbarium (Tucson, Arizona), and the Western Illinois University Fungarium (Macomb, Illinois).

Morphological characterization —

Mycelium from each isolate (TLT265, YP243, and AZ0501) was plated on seven types of media treated with tetracycline (50 mg/L) and streptomycin (50 mg/L). Hereafter, the use of antibiotics in media is represented with a plus sign “+”. The seven media tested were 1.3% malt extract agar (MEA+) (Difco Laboratories, Sparks, Maryland), Sabouraud dextrose agar (SDA+) (Oxoid LTD, Hampshire, England), Emerson yeast agar (EYSA+) (Difco Laboratories, Detroit, Michigan), Reasoner’s 2A agar (R2A+) (Difco Laboratories, Sparks, Maryland), Czapek-Dox Agar (CDA+) (Sigma-Aldrich, St. Louis, Missouri), peptone yeast agar (PYG+) (Mueller et al. 2004), and potato dextrose agar (PDA+) (Oxoid LTD, Hampshire, England). Media were prepared per manufacturers’ instructions. Cultures from each isolate were incubated separately at two temperatures, 25 °C and 35 °C, for 30 days in darkness. The Elcometer 6210 RAL Chart K7–

2014 was used as the color standard to describe the colonies. All cultures were produced from the same source culture for each strain.

Mycelial growth rates were measured on MEA+. Inoculum consisted of one 4 mm diameter agar plug from the growing edge of a fungal colony on MEA+. Plates were incubated at five different temperatures at 10 °C intervals from 5–45 °C in the dark. Plates were examined every five days for a month after cultures were established. Radial mycelial growth rates were calculated using arithmetic mean values per day by measuring colony diameter (Aoki et al. 2013). Three replicates were measured for each isolate at each temperature.

The production of sporangiopores was assessed using corn meal agar (CMA+) (Sigma-Aldrich, St. Louis, Missouri) supplemented with 50 mg/L chloramphenicol (Sigma-Aldrich, St. Louis, Missouri). Cultures were incubated at 25 °C and evaluated for 3 weeks.

Microscopy —

Tissue of each isolate was collected from the growing edge of a 15-day old colony on MEA+, SDA+, EYSA+, and PYG+ that had been incubated at 25 °C and 35 °C. Hyphae were stained with Bluemount (Hardy Diagnostics, Santa Maria, California) and compared under the light microscope.

Tissue was fixed for scanning electron microscopy (SEM) by incubation for 24 hours at 4°C in a fixative solution of 25 mM phosphate buffer (Sigma-Aldrich, St. Louis, Missouri) with a pH 7 and 3% gluteraldehyde (Sigma-Aldrich, St. Louis, Missouri). The fixative solution then was poured off and a 1% osmium tetroxide solution (Electron Microscopy Sciences, Hatfield, Pennsylvania) was added to the tissue. After incubating at 4 °C for 24 hours, the fungal tissue was rinsed three times with 25 mM phosphate buffer (pH 7) and then soaked in the following ethanol series for 15 minutes for dehydration of the tissue (30%, 50%, 65%, 75%, 89%, 95%,

3X-100%) and kept overnight at 100%, with two more 100% soaks the next day. After fixation, the specimens were dried in a carbon dioxide critical point dryer (Tousimis, Rockville, Maryland) and immediately placed in a sputter coater for a 2-minute coating of gold (Structure Probe, Inc., West Chester, Pennsylvania). The SEM images were obtained on a JEOL JSM-6010LA (JEOL Ltd. Tokyo, Japan) at the National Center for Agricultural Utilization Research, Peoria, Illinois, USA. Typical operating conditions were an accelerating voltage of 10kV and a spot size of 30.

DNA extraction, amplification, and sequencing —

DNA was extracted from the cultures obtained and 28S nuclear ribosomal RNA (nrRNA) was amplified following the protocols described below. The sequences were compared against GenBank (Altschul et al. 1997). The two cultures (TLT265 and YP243) matched the same phylotype as the one detected through culture-free sequencing (Hesse et al. 2016). The 28S nrRNA sequences for TLT265 and YP243 were compared against GenBank using BLASTn. One isolate with high similarity (>99%) from another study was identified (1125 BLAST score for both isolates; e value= 0). This isolate (AZ0501, accession number HM123225) was collected in 2007 in a study of endophytic fungi in a montane forest in southeastern Arizona (U'Ren et al. 2010). AZ0501 was isolated from photosynthetic tissue of a bryophyte (*Leucobryum* sp., Dicranaceae; U'Ren et al. 2010). This strain was obtained from the Robert L. Gilbertson Mycological Herbarium at The University of Arizona (accession number MYCO-ARIZ-AZ0501) and characterized.

Genomic DNA was extracted from cultures of each isolate using the Wizard genomic DNA purification kit (Promega, Madison, Wisconsin). PCR was used to amplify the partial 18S nrRNA with primers NS1 and NS4 (White et al. 1990), the partial 28S nrRNA using primers

LR0R and LR3 (Vilgalys and Hester 1990) and the ITS1-5.8S-ITS2 nrRNA (ITS barcode) using primers ITS1F (Gardes and Bruns 1993) and ITS4 (White et al. 1990). Each 25 µl of PCR mixture contained 12.5 µl of PCR master mix (Promega, Madison, Wisconsin), 3 µl of 1% bovine serum albumin (BSA), 1 µl of each primer (5 µM), 6.5 µl of nuclease-free water and 1 µl of DNA. Water was used in place of template for negative controls. The following PCR conditions were used: 95 °C for 5 min; 35 cycles of 94 °C for 30 s, annealing at 50 °C for 30 s, and extension at 72 °C for 45 s; and a final extension at 72 °C for 7 min. PCR products were checked using gel electrophoresis (1.2 % agarose in Tris-acetate-EDTA buffer). All samples were sequenced using the forward primer at Beckman Coulter Genomics (Danvers, Massachusetts). Sequences obtained were trimmed and edited using Sequencher (Gene Codes, Ann Arbor, Michigan). Similar sequences were identified by comparison with GenBank via the Basic Local Alignment Search Tool (BLASTn option) (Altschul et al. 1997) and comparison with the UNITE database (Kõljalg et al. 2013). The ITS-partial 28S nrRNA sequence of AZ0501 had been submitted previously to GenBank (HM123225; U'Ren et al. 2010). The sequences obtained here were deposited at GenBank (ITS rDNA, KU702608, KU702388; 28S rDNA, KU702551, KU702228; 18S rDNA, KU702505, KX372676, KX372677). The percentage similarity over the shared length for each locus among the three isolates (estimate of evolutionary divergence between sequences) was determined by the number of nucleotide differences per site between sequences in MEGA7 (Kumar et al. 2016).

Phylogenetic analyses —

Fungal sequences were edited in Geneious v.8.1.7 (<http://www.geneious.com>, Kearse et al. 2012), and used as queries to conduct BLAST searches for sequence similarity. Sequences were aligned with MUSCLE (Edgar et al. 2004). The alignments were trimmed with GBlocks

v.0.91b (Castresana 2000) using the less stringent conditions and then manually edited. The 18S and 28S rDNA datasets were concatenated into a single dataset of 1581 homologous nucleotide bases. Prior to phylogenetic inference, the best-fit models of nucleotide substitution were estimated using jModelTest v.2.1.9 (Darriba et al. 2012), for 18S and 28S rDNA data separately. Bayesian inference on the phylogeny of these taxa was carried out using MrBayes v.3.2.5 (Ronquist et al. 2012). The Markov chain Monte Carlo was run for 5 million generations under the TrN+I+G and TIM3+G nucleotide substitution models for 18S and 28S, respectively. Representatives of Mucoromycotina, Mortierellomycotina, and Glomeromycotina were selected to resolve the position of the novel fungal taxon within the recently erected phylum Mucoromycota (Spatafora et al. 2016). Alignments were submitted to TreeBASE (S19770).

Genome sequencing and phylogenomic analysis —

Isolate AZ0501 was grown in liquid yeast mold (YM) medium for approximately three weeks. Hyphal tissue was filtered from the growth medium and genomic DNA was extracted using the FastDNA SPIN Kit for soils (MP Biomedicals, Santa Ana, California). DNA was quantified using a QuBit. DNA was fragmented using a Covaris E220 and fragment sizing assessed using Agilent's BioAnalyzer. Fragments were made blunt and adenylated at the 3' end to ligate Illumina sequencing adapters. A unique index code was added during PCR amplification. Concentration was confirmed using KAPA's Library quantification kit on an ABI 7500 Real Time machine. The AZ0501 Illumina library was sequenced using Illumina's v2 500-cycle reagent kit as a spike-in with rRNA gene amplicon libraries on five separate Illumina MiSeq runs. On each run this isolate comprised about 30% of the sequences. Sequences from individual MiSeq runs were combined into a single FASTA file and assembled *de novo* using the Velvet software package (Zerbino 2010) with a k-mer value of 61 bp. Genome annotation for

developing genes for phylogenomic analyses was performed with MAKER (Holt and Yandell 2011) supported by *ab initio* gene prediction on the assembled contigs with Augustus (Stanke et al. 2004) using *Rhizopus delemar* parameters and self-trained GeneMarkHMM-ES (Ter-Hovhannisyan et al. 2008). Raw reads were deposited to the Sequence Read Archive (SRA) as Bioproject PRJNA368732 and Biosample SAMN06270744.

Phylogenomic analyses were performed with an approach previously applied for zygomycete genomes (Spatafora et al. in 2016). The data and analysis pipeline are available at https://github.com/zygolife/Bifiguratus_Phylogenomics. Briefly, the approach uses 434 protein coding genes that are typically single-copy in most fungi. Each phylogenetic marker is summarized as a profile Hidden Markov Model (HMM) (HMMER3, Eddy 2011) from an alignment of the orthologs from several dozen diverse fungi. The phylogenomic pipeline searches each HMM against the protein set of a species with HMMSEARCH and the best hit from each species is saved. A multiple sequence alignment (MSA) containing the orthologous copies from each species is constructed with HMMALIGN of these copies guided by the marker HMM. Each gene MSA is trimmed with TRIMAL (Capella-Gutiérrez et al. 2009) and the trimmed MSAs are concatenated into a single master alignment with a Perl script built with BioPerl (Stajich et al. 2002). Phylogenetic tree inference was performed with RAXML (v8.2.8) using autoMRE and rapid bootstrapping (-f a) (Stamatakis 2014).

Analysis of associated bacteria —

Mucoid growth was observed on the edges of cultures under some conditions, especially PYG+ at 35 °C. Light microscopy and culturing were used to evaluate the presence of bacteria on these mucoid areas. Bacteria present on fungal tissue were streaked onto R2A, from the three isolates grown on the seven media described above. The cultures were incubated at 25 °C for 7

days. We attempted to remove bacteria from the fungus using different antibiotics, media and temperatures with little success. However, their growth was especially limited on MEA+ at 25 °C.

DNA was extracted and amplified from the bacterial cultures following the protocol used for fungal samples (above). DNA amplification, sequencing and identification were performed as described above, except that we used universal primers for the 16S rRNA (8F and 1492R; Lane 1991) and an annealing temperature of 56 °C. Sequencing was performed using primer 1409R. Further sequencing of the *gyrA* gene of *Bacillus* strains was performed with primers p-*gyrA*-F and p-*gyrA*-R (Roberts et al. 1994) for identification (Chun and Bae 2000). Sequences were deposited to GenBank (KX397340–KX397345 for 16S; KX670823–KX670827 for *gyrA*).

Next generation sequencing was used with the purpose of confirming the presence of bacteria that may have evaded culturing. To assess dynamics of bacteria associated with this novel fungal taxon, isolates were grown on three media (PYG+, MEA+, and SDA+) at 25 °C and 35 °C. Three replicates were evaluated, one for each isolate (TLT265, YP243, and AZ0501), for a total of 18 samples. Tissue was collected from all samples and sent to MRDNA Molecular Research LP (Shallowater, Texas) for DNA extraction and Roche 454 FLX titanium sequencing. The 16S rRNA gene universal bacterial primer 27F (5'-AGRGTTTGATCMTGGCTCAG-3') was used in a modified version of amplicon pyrosequencing (bTEFAP®) (Dowd et al. 2008). A single-step 30 cycle PCR using HotStarTaq Plus Master Mix Kit (Qiagen, Valencia, California) was run under the following conditions: 94 °C for 3 minutes, followed by 28 cycles of 94 °C for 30 seconds; 53 °C for 40 seconds and 72 °C for 1 minute; with a final elongation step at 72 °C for 5 minutes. Amplicon products from different samples were mixed in equal concentrations and purified using Agencourt Ampure beads (Agencourt Bioscience Corporation, Beverly,

Massachusetts). Samples were sequenced with Roche 454 FLX titanium instruments and reagents following the manufacturer's procedures. With each new box of DNA extraction reagents two negative controls were run. No detectable amplification with universal 16S primers was detected on the controls.

Multiplex tags were used to parse the sequence library into 18 respective samples using the QIIME software package (Caporaso et al. 2010). The sequences obtained were denoised using AmpliconNoise (Quince et al. 2011), quality filtered and trimmed, and operational taxonomic units (OTUs) were clustered in USEARCH (Edgar 2010) using the UPARSE (Edgar 2013) algorithm considering a pairwise sequence identity percentage of 0.97. Chimera filtering was performed using UCHIME (Edgar et al. 2011) comparing to the RDP Gold database (http://drive5.com/usearch/manual/cmd_uchime_ref.html). Taxonomy was assigned using the RDP Classifier (Wang et al. 2007) against the Greengenes database (DeSantis et al. 2006) and corroborated with BLAST (Altschul et al. 1997) matches against GenBank (Benson et al. 2011). Singletons and sequences assigned to mitochondria were removed. Raw reads were deposited to the Sequence Read Archive (SRA) as study accession SRP082174 (Bioproject PRJNA318907).

RESULTS

Two isolates of the focal fungus (YP243 and TLT265) were obtained using filtration-pasteurization methods from soil. Both isolates were obtained from the lower A horizon (5–10 cm deep) of a N-fertilized plot under ambient CO₂ in a pine plantation in North Carolina, USA. Isolates YP243 and TLT265 were obtained from the 63 mm and 2000 mm particle size, respectively, and grown at 25 °C on PYG+ (Hesse et al. 2016). These isolates were evaluated with the only known, cultured closest relative in GenBank, isolate AZ0501, which was obtained from surface-sterilized photosynthetic tissue of a moss (U'Ren et al. 2010). Isolates YP243 and

TLT265 share >99% similarity for the ITS, 18S and 28S genes sequences with AZ0501 (TABLE S1). Morphological characterization of the isolates is provided in the taxonomic description, including growth on different media (FIGS. 1–2) and microscopy (FIGS. 3–4).

Phylogenetic analyses —

Given the lack of available data for genomes of fungi such as *Endogone* and *Sphaerocreas*, and because ITS nrRNA sequences could not be aligned across this phylum, concatenated 18S and 28S nrRNA analyses containing more complete taxon sampling were used to estimate the phylogenetic placement of these strains within Mucoromycotina. Isolates AZ0501, TLT265 and YP243 clustered together within a well-supported monophyletic clade that branched after early-diverging lineages such as *Calcarisporiella* (Hirose et al. 2012), *Densospora* (McGee 1996), *Endogone* (Yamamoto et al. 2015) and *Sphaerocreas* (Hirose et al. 2014) and unidentified Mucoromycotina (Truong et al. 2017) (FIG. 5).

Genome sequencing and phylogenomic analysis —

Sequencing generated a total of approx. 6.8M high quality reads, of which 5.6M reads assembled into 568 scaffolds of at least 500bp. The final assembly had an N50 of ~928kbp, an overall genome assembly length of ~19.5 Mbp, and a GC content of 48.3 %. The taxon represented by these three strains has 8876 predicted protein-coding genes. This is a nearly complete genome, but lacks RNA transcriptome data to provide additional support for gene models. The assembly shows a high degree of completeness with a Benchmarking Universal Single-Copy Orthologs (BUSCO) score of 82.1% using Fungi ODB9 set. The genome is 86% complete. Out of the 290 BUSCO groups searched, we found 217 Complete Single-Copy, 21 Complete Duplicated, 12 Fragmented, and 40 missing BUSCO orthologs (Simão et al. 2015). These results are similar to other Mucoromycotina (TABLE 1). Indicative of a nearly complete

genome assembly, we identified 237 of 248 super-conserved Core Eukaryotic Genes (CEGs). The CEGMA pipeline (Parra et al. 2007) suggests 95.6% genome completeness (TABLE S2).

The Maximum Likelihood phylogenetic tree generated with RAXML was produced from a concatenated alignment of 434 protein-coding genes (165,307 residues across ten taxa). The phylogenomic tree places the new taxon as one of the early diverging lineages in Mucoromycotina, only preceded by *Calcarisporiella*, the first diverging known lineage in the subphylum (FIG. 6).

Analysis of associated bacteria —

Two genera of bacteria were successfully isolated and identified based on 16S rRNA data: *Bacillus licheniformis* (KX397340–KX397344) and *Stenotrophomonas* sp. (KX397345). Due to the limitations of the 16S rRNA gene in resolving members of the *Bacillus subtilis* complex, identities of the *Bacillus licheniformis* strains were confirmed using the *gyrA* gene (KX670823–KX670827) (TABLE S3). Both bacteria were recovered from PYG+, SDA+, and CZP+ at 25 °C. Although bacteria were not isolated from fungal isolate AZ0501, they were observed on this isolate using SEM (FIG. 7C, 7D), and were confirmed by direct sequencing (FIG. 7A).

Culture-independent methods based on 454 amplicon pyrosequencing were used to assess bacterial communities associated with the three isolates grown on different media at two temperatures (25 °C and 35 °C). A majority of the sequences turned out to be fungal mitochondria amplicons, which were removed from the analysis. The remaining data show numerous bacterial taxa associated with these three fungal isolates (Fig. 7A). However, aside from *Bacillus*, *Methylobacterium* spp. and *Propionibacterium* most bacterial taxa were in low read abundance (<10 sequences / OTU), and in four of the 18 treatments no bacteria were

detected (TABLE S4). Venn diagrams (Fig. 7B) show that few bacterial OTUs are shared between isolates, with the majority of OTUs being distinct for each isolate. *Bacillus* and *Methylobacterium* were the dominant taxa associated with YP243 and TLT265, however *Bacillus* was not detected from isolate AZ0501 in the 454 dataset, nor when isolates were grown on MEA+ and SDA+ media (FIG. 7A, 7B). Isolates TLT265 and YP243 show a higher bacterial richness with respect to isolate AZ0501. Variation in bacterial taxa and their relative abundance (in terms of sequence reads) was observed among temperature and media treatments (FIG. 7A, 7B). A greater number of bacterial OTUs were detected from strains obtained from soil and those growing on MEA+ (FIG. 7B). Although a core set of bacterial OTUs are present at both temperatures (25 °C and 35 °C), incubation temperature appeared to shift the bacterial communities in these isolates. In contrast, three *Methylobacterium* OTUs were consistently detected from all three isolates at 25 °C and 35 °C on MEA+ and SDA+.

TAXONOMY

Bifiguratus T. J. Torres-Cruz & A. Porras-Alfaro, gen. nov.

MycoBank MB 818093

FIGS. 1, 3–4.

Typification: *Bifiguratus adalaidae* T. J. Torres-Cruz and A. Porras-Alfaro.

Etymology: from the Latin “*figuratus*”, in reference to the dimorphic characteristic of the type species.

Bifiguratus adalaidae T. J. Torres-Cruz & A. Porras-Alfaro, sp. nov.

MycoBank MB 818094

FIGS. 1, 3–4.

Typification: UNITED STATES: Arizona, Cochise County, Coronado National Forest,

eastern Chiricahua Mountains, dried culture from an isolate from surface-sterilized
photosynthetic tissue of *Leucobryum* sp., Arizona, USA, Jun 2007, *J. M. U'Ren, A. E. Arnold, et*
al. (**holotype** MICH 134798). Ex-type culture MYCO–ARIZ–AZ0501 = NRRL 66559.

Etymology: In honor of Adelaida Chaverri Polini, a female Costa Rican scientist.

Full description —

Cultures of the three strains grown on MEA+ at 25 °C present a white (RAL 9003)
smooth flat colony with entire margins (FIG. 1A). At 35 °C colonies are smaller than at 25 °C,
white (RAL 9003), and 1–2 mm raised with undulate margins and a rough rugose surface (FIG.
1E). On SDA+ at 25 °C colonies are cream (RAL 9001), rough, and 1–2 mm raised with entire
margins (FIG. 1B), while at 35 °C the colonies have the same characteristics but are more
corrugated (FIG. 1F). Colony characteristics on CMA+ at 25 °C are consistent with those
observed on SDA+. On EYSA+ at 25 °C colonies are pure white (RAL 9010), rough, 2–3 mm
high colonies have lobate margins (FIG. 1C). On this medium at 35 °C, colonies are light ivory
(RAL 1015) and flat but still wrinkled, with a filamentous margin (FIG. 1G) and a more mucoid
texture is observed at the edges with a cream pigmentation (RAL 9001). On PYG+ at 25 °C
colonies are highly corrugated (FIG. 1D), irregular, cream (RAL 9001), with undulated margins,
and are 3–4 mm high. At 35 °C on PYG+ colonies are cream (RAL 1001), corrugated, and
raised, but the edges present a mucoid growth with undulated margins (FIG. 1H) that have a
translucent beige-red color (RAL 3012). The presence of a mucoid texture at the edges of the
fungus growing at 35 °C on EYSA+ and PYG+ (FIGS. 1G–H) is related to more abundant growth
of bacteria in the cultures, with respect to MEA+ and PDA+. On PDA+ the fungus has the same
morphological characteristics as those on PYG+ at 25 °C, and the morphology remains the same
at 35 °C. None of the strains grew on R2A and CDA; fungal-associated bacteria colonized the

plates instead. All cultures were produced from the same source culture for each strain, such that the bacteria appear to have been present in the source and became dominant on R2A and CDA media.

As cultures aged (i.e., after more than one month of incubation) the morphology of the cultures changed. In some cases (e.g., SDA+), colonies became more mucoid and flat, and were overcome by bacterial growth. However, in most cases (e.g. MEA+), cultures presented more hyphal growth at the edges and colonies became darker, turning brown.

Colonies on MEA+ have radial mycelial growth rates of 0.4 mm/d at 25 °C and 0.2 mm/d at 35 °C in the dark, during the first 15 days of incubation. Growth rates at different incubation temperatures are shown in FIG. 2. The three isolates did not grow at 5 °C or at 45 °C. The isolates generally grew more rapidly at 25 °C during the first two weeks of incubation. After 15 days of growth, the growth rate decreased at 25 °C and increased at 35 °C for YP243 and AZ0501.

The three strains possess coenocytic mycelium with hyphae ca. 2 µm in diameter (15-day old culture, MEA+, 25 °C; FIGS. 3B and 4B). When observed by SEM, hyphal surfaces present rough texture with surface ornamentations (FIG. 4B–C). No sporangiospores were observed, although multiple attempts were made using PDA, CMA, and several temperatures. The fungus produces what appear to be intercalary and terminal chlamydospores 5–10 µm in size, which appear globose to sub-globose (FIGS. 3B, 4A), but additional studies are necessary to confirm the nature of these structures. SEM observations revealed the presence of yeast-form cells (FIGS. 4D–F). Bud scars and budding yeasts were observed (FIGS. 4D–E), and germ tubes were visualized on yeast-form cells (FIG. 4F). They were commonly observed when the fungus was cultivated on MEA+ (FIG. 4E) and EYSA+ (FIGS. 4D, F) at 25 °C. Thus these isolates appear to be dimorphic.

Other cultures examined: UNITED STATES, North Carolina, Blackwood Division of the Duke Forest free air CO₂ enrichment (FACE) experiment (35°58'N, 79°06'W; elevation 130 m), Oct 2013, *C. R. Kuske*, MICH 134800 (NRRL 66560 = WIU–APA–TLT265) and MICH 134801 (NRRL 66561 = WIU–APA–YP243).

Substrate —

Because only three cultures are currently available for this fungus, we based analyses of substrates and potential hosts on analyses of available ITS rRNA data from BLASTn analysis. Two cultures (YP243 and TLT265) were recovered from soil, and apparently related fungi have been detected on several other soil samples using direct sequencing from environmental samples including temperate forest soils (HQ021892, DQ421188, DQ421187), truffle grounds soils (JF927042, JF927041), rice field soils (AB520601), and humus under *Pinus sylvestris* (GU559100). See ‘distribution’ section for geographic range information.

Our previous report suggested that this species represented up to 9% of the total 454 sequences in individual samples at the Duke Forest (Hesse et al. 2016). It is rarely found in the forest floor as is more common on the deeper horizons (Hesse et al. 2016). Hesse et al. (2016) showed that the species increases in abundance with soil depth and reaches the highest abundance in a 5–10 cm depth. It responds to the combination of N fertilization and CO₂ by increasing in relative abundance at the 2–5 cm and 5–10 cm deep intervals (Hesse et al. 2016).

In addition to occurring as a culturable endophyte in *Leucobryum* (AZ0501, U'Ren et al. 2010), environmental data show that related strains have been detected as a root associated fungus of the orchid *Cymbidium kanran* (KU141166, Hong et al. 2015) and sweet chestnut, *Castanea sativa* (EF040836, Peintner et al. 2007).

Distribution —

This taxon is known from the Northern Hemisphere. In the United States, phylotypes consistent with *B. adelaidae* have been detected from the Duke Forest, North Carolina (YP243 and TLT265), Bartlett Experimental Forest, New Hampshire (HQ021892), Cedar Creek Ecosystem Science Reserve, Minnesota (DQ421188, DQ421187), and Coronado National Forest, Arizona (AZ0501). It has also been detected in Italy (JF927042, JF927041, EF040836), Japan (AB520601), Sweden (GU559100), and South Korea (KU141166).

DISCUSSION

Fungal isolates obtained from soil (TLT265, YP243) and as an endophyte (AZ0501) provided the opportunity for morphological, multi-locus and phylogenomic description of a new endophytic and soil-dwelling species of Mucoromycota with an apparent wide distribution across the Northern Hemisphere. This dimorphic species is affiliated with epihyphal bacteria and appears to have a wide host range spanning multiple lineages of plants, as well as subsurface soils in diverse forests. This fungus is part of UNITE's "most wanted" fungi as cluster code UCL7_006587, ranked #1 at the phylum level based on number of studies and #4 based on number of sequences (Nilsson et al. 2016). Description of this species exemplifies the potential to use environmental sequencing to guide taxonomic discovery. Its description and characterization represents an effort to help bridge the gap between fungal taxonomy and molecular ecology.

Asexual reproduction among lineages formerly known as zygomycetes can be achieved by the formation of chlamydospores, sporangiospores, yeast cells, or arthrospores (Fuller 1978; Benny et al. 2014). *Bifiguratus adelaidae* presents coenocytic hyphae, globose intercalary chlamydospores, and yeast cells. The presence of yeast cells in other Mucoromycotina taxa has been described (e.g., *Mucor*, *Benjaminiella* and *Mycotypha*; Benny et al. 2014). After budding,

scar tissue is left behind and the mother cell will exhibit a crater-like ring of scar tissue (Chant and Pringle 1995; Powell et al. 2003), consistent with those observed in *B. adalaidae* (FIG. 4D). Budding yeasts were also viewed (FIG. 4E). Chlamydospores form in swellings of coenocytic young hyphae (Benny et al. 2014). We observed what appears to be germinating chlamydospores (FIG. 4F), which also have been observed in other dimorphic fungi (e.g., Ascomycota such as *Candida albicans* and *Candida dubliniensis*). In those species, chlamydospores can generate daughter chlamydospores, bud to form daughter yeast cells, and germinate to produce pseudohyphae and hyphae (Citiulo et al. 2009). Chlamydospores have been reported in other Mucoromycotina fungi, including *Umbelopsis* (Wang et al. 2013), *Ambomucor* (Liu and Zheng 2015), and *Isomucor* (Ivanildo et al. 2012). Chlamydospore formation can be induced by bacteria associated with fungi, as seen in three Mucoromycotina species (*Phycomyces blakesleeanus*, *Mucor bacilliformis* and *Mucor hiemalis*) where a lipopeptide produced by *Ralstonia solanacearum* induces the production of chlamydospores as a survival technique for the fungus (Spraker et al. 2016). It is possible that the presence of bacteria on *B. adalaidae* could influence chlamydospore production (FIG. 3B), a topic that will require further study.

Phylogenetic inferences based on the concatenated partial 18S and 28S rRNA loci placed *B. adalaidae* as a novel taxon in Mucoromycota (FIG. 5). The *B. adalaidae* branch diverges after *Calcarisporiella*, the earliest known branching Mucoralean fungus or sister to Mucoromycotina. It is distinct from taxa such as *Endogone*, *Sphaerocreas*, and *Umbelopsis* (FIG. 6).

Fungi in Mucoromycotina and related lineages in Mucoromycota often associate with other organisms, including plants and bacteria (Benny et al. 2014, Bonfante and Desirò 2017). Glomeromycotina form arbuscular mycorrhizal associations, some species of Mortierellomycotina can associate with mosses and other plants (Melo et al. 2014), and diverse

Mucoromycotina present associations with living plants. For example, some Mucoromycotina associate with liverworts in the thallus midrib and subterranean axes (Field et al. 2015a; Field et al. 2015b). Species of *Umbelopsis* associate with roots of orchids (Yu et al. 2015), and some *Mucor* and *Umbelopsis* have been isolated from woody roots of ponderosa pine and Douglas fir (Hoff et al. 2004). In this study, isolates of the new lineage *B. adelaidae* were obtained from soil and as a putative endophyte of *Leucobryum* sp. (U'Ren et al. 2010). Additional studies are needed to assess whether these fungi function in a symbiotic capacity with plants.

Many lineages of early diverging terrestrial fungi often associate with bacteria. For example, Gram-positive endobacteria related to the Mollicutes occur in cytoplasm of Glomeromycotina (Naumann et al. 2010; Toomer et al. 2015). Recent studies have provided evidence of the presence of bacteria in some Mucoromycotina, such as *Endogone*, where similar Mollicutes-related endobacteria have been detected (Desirò et al. 2015). Endobacteria occur in the cytoplasm of Mortierellomycotina, like *Mortierella elongata* (Sato et al. 2010; Uehling et al. 2017) and *Rhizopus* (Partida-Martinez and Hertweck 2005). Based on SEM micrographs we conclude that at least some of these bacteria are epiphythal in *B. adelaidae*, but additional testing is necessary to determine if part of the taxa detected on this fungus are endobiotic. To our knowledge this is the first report of *Bacillus licheniformis*, *Methylobacterium* and *Stenotrophomonas* living in association with a fungus (but see the *Bacillus* clade identified by Hoffman and Arnold 2010).

ACKNOWLEDGMENTS

This study was supported through a Science Focus Area grant to CRK from the U.S. Department of Energy, Biological and Environmental Research Division. TJTC thanks the RISE (Research Inspiring Student Excellence) and Women in Science programs at WIU for research

support. GMNB, AD and GB thank MSU AgBioResearch and USDA for research support. Isolation and sequencing of AZ0501 were supported by the efforts of Jana U'Ren, Jolanta Miadlikowska, François Lutzoni, Alex Laetsch, Fabiola Santos, and Ester Gaya, with financial support from the College of Agriculture and Life Sciences at The University of Arizona, NSF DEB-0640996 to AEA, and DEB-0640956 to François Lutzoni. JES was supported by NSF DEB-1441715 and high performance computing resources at the Institute for Integrative Genome Biology at University of California-Riverside supported by NSF DBI-1429826 and NIH S10-OD016290. The authors thank Arthur Thompson (National Center for Agricultural Utilization Research, Peoria, IL) for generating the SEM images and Adrian Tsang (Concordia University) for access to the *Calcarisporiella thermophila* genome. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture or National Science Foundation. The mention of firm names or trade products does not imply that they are endorsed or recommended by the USDA over other firms or similar products not mentioned. USDA is an equal opportunity provider and employer.

The type species described here was named in honor of Adelaida Chaverri Polini. Dr. Chaverri was a world-recognized tropical biologist and role model for women in Latin America in science, technology, engineering, and mathematics. Her major contributions were in the study and conservation of tropical montane forests and treeless alpine grasslands (*páramos*). As a biologist, she recognized the importance of mycorrhizal fungi and contributed to the description of a fungus in the Acaulosporaceae, a fungus in the same phylum as the species described here (Sieverding et al. 1988). She supported the creation of Costa Rica's National Park Service, providing the basis for extensive conservation of biological diversity. In 2013, she was included

in the *Galería de la Mujer* (Women Gallery), one of the highest honors given to Costa Rican women for their dedication to defending and improving women human rights (Kappelle and Cleef 2004).

LITERATURE CITED

- Altschul SF, Madden T, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman D J. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25:3389–3402.
- Aoki T, Smith JA, Mount LL, Geiser DM, O'Donnell K. 2013. *Fusarium torreyae* sp. nov., a pathogen causing canker disease of Florida torreyia (*Torreya taxifolia*), a critically endangered conifer restricted to northern Florida and southwestern Georgia. *Mycologia* 105:312–319.
- Benny GL, Humber RA, Voigt K. 2014. Zygomycetous fungi: Phylum entomophthoromycota and subphyla Kickxellomycotina, Mortierellomycotina, Mucoromycotina, and Zoopagomycotina. In: McLaughlin DJ, Spatafora JW, eds. *Systematics and evolution*. New York: Springer-Verlag. p 209–239.
- Benson AD, Karsch-Mizrachi I, Clark K, Lipman DJ, Ostell J, Sayers EW. 2011. GenBank. *Nucleic Acids Research* 26:1–7.
- Blackwell, M. 2011. The Fungi: 1, 2, 3... 5.1 million species? *American Journal of Botany* 98:426–438.
- Bonfante P, Desirò A. 2017. Who lives in a fungus? The diversity, origins and functions of fungal endobacteria living in Mucoromycota. *ISME Journal* doi:10.1038/ismej.2017.21
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. TrimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973.

- 532 Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N,
 533 Gonzalez Pena A, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE,
 534 Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR,
 535 Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010. QIIME
 536 allows analysis of high-throughput community sequencing data. *Nature Methods* 7:335–336.
- 537 Castresana, J. 2000. Selection of conserved blocks from multiple alignments for their use in
 538 phylogenetic analysis. *Molecular Biology and Evolution* 17:540–552.
- 539 Chant J, Pringle JR. 1995. Patterns of bud-site selection in the yeast *Saccharomyces cerevisiae*.
 540 *Journal of Cell Biology* 129:751–765.
- 541 Chun J, Bae KS. 2000. Phylogenetic analysis of *Bacillus subtilis* and related taxa based on partial
 542 *gyrA* gene sequences. *Antonie van Leeuwenhoek* 78:123–127.
- 543 Citiulo F, Moran GP, Coleman DC, Sullivan DJ. 2009. Purification and germination of *Candida*
 544 *albicans* and *Candida dubliniensis* chlamydospores cultured in liquid media. *FEMS Yeast*
 545 *Research* 9:1051–1060.
- 546 Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics
 547 and parallel computing. *Nature Methods* 9:772.
- 548 DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,
 549 Andersen G.L. 2006. Greengenes, a chimera-checked 16S rRNA gene database and
 550 workbench compatible with ARB. *Applied and Environmental Microbiology* 72:5069–5072.
- 551 Desirò A, Faccio A, Kaech A, Bidartondo MI, Bonfante P. 2015. *Endogone*, one of the oldest
 552 plant-associated fungi, host unique Mollicutes-related endobacteria. *New Phytologist*
 553 205:1464–1472.

- 554 Dowd SE, Sun Y, Wolcott RD, Doming A, Carroll JA. 2008. Bacterial tag-encoded FLX
 555 amplicon pyrosequencing (bTEFAP) for microbiome studies: bacterial diversity in the ileum
 556 of newly weaned *Salmonella*-infected pigs. Foodborne Pathogens and Disease 5:459–472.
- 557 Eddy SR. 2011. Accelerated Profile HMM Searches. PLoS Comput Biol 7:e1002195.
- 558 Edgar RC. 2014. MUSCLE: multiple sequence alignment with high accuracy and high
 559 throughput. Nucleic acids research 32:1792–1797.
- 560 Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics
 561 26:2460–2461.
- 562 Edgar RC. 2013. UPARSE: Highly accurate OTU sequences from microbial amplicon reads.
 563 Nature Methods 10:996–998.
- 564 Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and
 565 speed of chimera detection. Bioinformatics 27:2194–2200.
- 566 Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, Duckett JG,
 567 Leake JR, Pressel S. 2015a. First evidence of mutualism between ancient plant lineages
 568 (Haplomitriopsida liverworts) and Mucoromycotina fungi and its response to simulated
 569 Palaeozoic changes in atmospheric CO₂. New Phytologist 205:743–756.
- 570 Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, Duckett JD,
 571 Leake JR, and Pressel S. 2015b. Functional analysis of liverworts in dual symbiosis with
 572 Glomeromycota and Mucoromycotina fungi under a simulated Palaeozoic CO₂ decline.
 573 ISME Journal 10:1514–1526.
- 574 Fuller MS. 1978. Lower fungi in the laboratory. Georgia: Department of Botany, University of
 575 Georgia. p 213.

- 576 Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes
577 application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2:113–118.
- 578 Grigoriev IV, Cullen D, Goodwin SB, Hibbett D, Jeffries TW, Kubicek CP, Kuske C, Magnuson
579 JK, Martin F, Spatafora JW, Tsang A, Baker SE. 2011. Fueling the future with fungal
580 genomics. *Mycology* 2:192–209.
- 581 Gupta VK, Tuohy MG, Ayyachamy M, Turner KM, O'Donovan A. 2012. Laboratory protocols
582 in fungal biology: current methods in fungal biology. New York: Springer-Verlag. p 69–70.
- 583 Hesse CN, Torres-Cruz TJ, Tobias TB, Al-Matruk M, Porras-Alfaro A, Kuske CR. 2016.
584 Detection and targeted isolation of novel nitrogen-responsive fungal taxa from temperate
585 pine forest soil. *Mycologia* 108:1049–1068.
- 586 Hirose D, Degawa Y, Inaba S, Tokumasu S. 2012. The anamorphic genus *Calcarisporiella* is a
587 new member of Mucoromycotina. *Mycoscience* 53:256–260.
- 588 Hirose D, Degawa Y, Yamamoto K, Yamada A. 2014. *Sphaerocreas pubescens* is a member of
589 the Mucoromycotina closely related to fungi associated with liverworts and hornworts.
590 *Mycoscience* 55:221–226.
- 591 Hoff JA, Klopfenstein NB, McDonald GI, Tonn JR, Kim MS, Zambino PJ, Hessburg PF, Rogers
592 JD, Peever TL, Carris LM. 2004. Fungal endophytes in woody roots of Douglas-fir
593 (*Pseudotsuga menziesii*) and ponderosa pine (*Pinus ponderosa*). *Forest Pathology* 34:255–
594 271.
- 595 Hoffman MT, Arnold AE. 2010. Diverse bacteria inhabit living hyphae of phylogenetically
596 diverse fungal endophytes. *Applied Environmental Microbiology* 76:4063–4075.
- 597 Holt C, Yandell M. 2011. MAKER2: an annotation pipeline and genome-database management
598 tool for second-generation genome projects. *BMC Bioinformatics* 12:491.

- 599 Hong JW, Suh H, Kim OH, Lee NS. 2015. Molecular identification of mycorrhizae of
 600 *Cymbidium kanran* (Orchidaceae) on Jeju Island, Korea. *Mycobiology* 43:475–480.
- 601 Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,
 602 Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson
 603 R, Thomas BC, Banfield JF. 2016. A new view of the tree of life. *Nature Microbiology*
 604 1:16048.
- 605 Ivanildo de Souza JI, Amorim CL, dos Santos JF, Pinheiro J, Harakava R. 2012. *Isomucor*
 606 (Mucoromycotina): a new genus from a Cerrado reserve in state of Sao Paulo, Brazil.
 607 *Mycologia* 104:232–241.
- 608 Kappelle M, Cleef AM. 2004. Adelaida Chaverri-Polini. *Revista de Biología Tropical* 52:XIII-
 609 XVI.
- 610 Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
 611 Markowitz S, Duran C, Thierer T, Ashton B, Mentjies P, Drummond A. 2012. Geneious
 612 Basic: an integrated and extendable desktop software platform for the organization and
 613 analysis of sequence data. *Bioinformatics* 28:1647–1649.
- 614 Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates ST, Bruns
 615 TD, Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Dueñas M,
 616 Grebenc T, Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lücking
 617 R, Martín MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson
 618 M, Põldmaa K, Saag L, Saar I, Schüßler A, Scott JA, Senés C, Smith ME, Suija A, Taylor
 619 DL, Telleria MT, Weiß M, Larsson K-H. 2013. Towards a unified paradigm for sequence-
 620 based identification of Fungi. *Molecular Ecology* 22:5271–5277.

- 621 Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis
622 version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33:1870–1874.
- 623 Lane DJ. 1991. 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M, eds. *Nucleic acid*
624 *techniques in bacterial systematics*. New York: Wiley. p 115–175.
- 625 Liu XY, Zheng RY. 2015. New taxa of *Ambomucor* (Mucorales, Mucoromycotina) from China.
626 *Mycotaxon* 130:165–171.
- 627 Martin F, Cullen D, Hibbett D, Pisabarro D, Spatafora JW, Baker SE, Grigoriev IV. 2011.
628 Sequencing the fungal tree of life. *New Phytologist* 190:818–821.
- 629 McGee PA. 1996. The Australian zygomycetous mycorrhizal fungi: the genus *Densospora* gen.
630 nov. *Australian Systematic Botany* 9: 329–336.
- 631 McMurdie PJ, Holmes S. 2013. phyloseq: An R package for reproducible interactive analysis
632 and graphics of microbiome census data. *PLoS ONE* 8:e61217.
- 633 Melo IS, Santos SN, Rosa LH, Parma MM, Silva LJ, Queiroz SCN, Pellizari VH. 2014.
634 *Mortierella alpina* strain from the Antarctic moss *Schistidium antarctici*. *Extremophiles*
635 18:15–23.
- 636 Mueller G, Bills G, Foster M. 2004. Biodiversity of fungi: inventory and monitoring methods.
637 San Diego, California: Elsevier Academic Press. 777p.
- 638 Naumann M, Schüßler A, Bonfante P. 2010. The obligate endobacteria of arbuscular mycorrhizal
639 fungi are ancient heritable components related to the Mollicutes. *ISME Journal* 4:862–871.
- 640 Nilsson RH, Wurzbacher C, Bahram M, Coimbra VRM, Larsson E, Tedersoo L, Eriksson J,
641 Duarte Ritter C, Svantesson S, Sánchez-García M, Ryberg M, Kristiansson E, Abarenkov K.
642 2016. Top 50 most wanted fungi. *MycoKeys* 12:29–40.

- 643 Parra G, Bradnam K, Korf I. 2007. CEGMA: a pipeline to accurately annotate core genes in
644 eukaryotic genomes. *Bioinformatics* 23:1061–1067.
- 645 Partida-Martinez L, Hertweck C. 2005. Pathogenic fungus harbours endosymbiotic bacteria for
646 toxin production. *Nature* 437:884–888.
- 647 Peintner U, Lotti M, Klotz P, Bonuso E, Zambonelli A. 2007. Soil fungal communities in a
648 *Castanea sativa* (chestnut) forest producing large quantities of *Boletus edulis sensu lato*
649 (porcini): where is the mycelium of porcini? *Environmental Microbiology* 9:880-889.
- 650 Powell CD, Quain DE, Smart KA. 2003. Chitin scar breaks in aged *Saccharomyces cerevisiae*.
651 *Microbiology* 149:3129–3137.
- 652 Quince C, Lanzen A, Davenport RJ, Turnbaugh PJ. 2011. Removing noise from pyrosequenced
653 amplicons. *BMC Bioinformatics* 12:38.
- 654 Roberts MS, Nakamura LK, Cohan FM. 1994. *Bacillus mojavensis* sp. nov., distinguishable from
655 *Bacillus subtilis* by sexual isolation, divergence in DNA sequence, and differences in fatty
656 acid composition. *International Journal of Systematic Bacteriology* 44:256–264.
- 657 Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,
658 Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference
659 and model choice across a large model space. *Systematic Biology* 61:539–542.
- 660 Sato Y, Narisawa K, Tsuruta K, Umezu M, Nishizawa T, Tanaka K, Yamaguchi K, Komatsuzaki
661 M, Ohta H. 2010. Detection of betaproteobacteria inside the mycelium of the fungus
662 *Mortierella elongata*. *Microbes and Environments* 25:321–324.
- 663 Sieverding E, Chaverri A, Rojas I. 1988. *Acaulospora splendida*, a new species in the
664 Endogonaceae from Costa Rica. *Mycotaxon* 33:251–256.

- 665 Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
666 assessing genome assembly and annotation completeness with single-copy orthologs.
667 Bioinformatics 31:3210–3212.
- 668 Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G, Corradi N,
669 Grigoriev I, Gryganskyi A, James TY, O'Donnell K, Taylor TN, Uehling J, Vilgalys R,
670 White M, Stajich JE. 2016. Zygomycete Genealogy of Life (ZyGoLife): a phylum-level
671 phylogenetic classification of zygomycete fungi based on genome-scale data. Mycologia
672 108:1028–1046.
- 673 Spraker JE, Sanchez LM, Lowe TM, Dorrestein PC, Keller NP. 2016. *Ralstonia solanacearum*
674 lipopeptide induces chlamydospore development in fungi and facilitates bacterial entry into
675 fungal tissues. ISME Journal 10:2317–2330.
- 676 Stajich JE, Block D, Boulez K, Brenner SE, Chervitz SA, Dagdigian C, Fuellen G, Gilbert JG,
677 Korf I, Lapp H, Lehväslaiho H, Matsalla C, Mungall CJ, Osborne BI, Pocock MR, Schattner
678 P, Senger M, Stein LD, Stupka E, Wilkinson MD, Birney E. 2002. The Bioperl toolkit: Perl
679 modules for the life sciences. Genome Research 12:1611–1618.
- 680 Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of
681 large phylogenies. Bioinformatics 30:1312–1313.
- 682 Stanke M, Steinkamp R, Waack S, Morgenstern B. 2004. AUGUSTUS: a web server for gene
683 finding in eukaryotes. Nucleic Acids Research 32(suppl 2):W309–W312.
- 684 Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008. Gene prediction in
685 novel fungal genomes using an *ab initio* algorithm with unsupervised training. Genome
686 Research 18:1979–1990.

- 687 Toomer KH, Chen X, Naito M, Mondo SJ, Bakker HC, VanKuren NW, Lekberg Y, Morton JB,
 688 Pawlowska TE. 2015. Molecular evolution patterns reveal life history features of
 689 mycoplasma-related endobacteria associated with arbuscular mycorrhizal fungi. *Molecular*
 690 *Ecology* 24:3485–3500.
- 691 Truong C, Mujic AB, Healy R, Kuhar F, Furci G, Torres D, Niskanen T, Sandoval-Leiva PA,
 692 Fernández N, Escobar JM, Moretto A, Palfner G, Pfister D, Nouhra E, Swenie R, Sánchez-
 693 García M, Matheny PB, Smith M. 2017. How to know the fungi: combining field inventories
 694 and DNA-barcoding to document fungal diversity. *New Phytologist* 214:913–919.
- 695 U'Ren JM, Lutzoni F, Miadlikowska J, Arnold AE. 2010. Community analysis reveals close
 696 affinities between endophytic and endolichenic fungi in mosses and lichens. *Microbial*
 697 *Ecology* 60:340–353.
- 698 Uehling J, Gryganskyi A, Hameed K, Tschaplinski T, Misztal PK, Wu S, Desirò A, Vande Pol
 699 N, Du Z, Zienkiewicz A, Zienkiewicz K, Morin E, Tisserant E, Splivallo R, Hainaut M,
 700 Henrissat B, Ohm R, Kuo A, Yan J, Lipzen A, Nolan M, LaButti K, Barry K, Goldstein AH,
 701 Labbé J, Schadt C, Tuskan G, Grigoriev I, Martin F, Vilgalys R, Bonito G. 2017.
 702 Comparative genomics of *Mortierella elongata* and its bacterial endosymbiont *Mycoavidus*
 703 *cysteinexigens*. *Environmental Microbiology*. doi:10.1111/1462-2920.13669
- 704 Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping enzymatically amplified
 705 ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172:4238–4246.
- 706 Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naïve Bayesian classifier for rapid assignment
 707 of rRNA sequences into the new bacterial taxonomy. *Applied Environmental Microbiology*
 708 73:5261–5267.

- Wang Y, Liu X, Zheng R. 2013. Four new species records of *Umbelopsis* (Mucoromycotina) from China. *Journal of Mycology* 2013:1–6.
- Weber CF, Vilgalys R, Kuske CR. 2013. Changes in fungal community composition in response to elevated atmospheric CO₂ and nitrogen fertilization varies with soil horizon. *Frontiers in Microbiology* 4:78.
- White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White, TJ, eds. *PCR protocols a guide to methods and applications*. New York: Academic Press. p 315–322.
- Yamamoto K, Degawa Y, Hirose D, Yamada A. 2015. Morphology and phylogeny of four *Endogone* species and *Sphaerocreas pubescens* collected in Japan. *Mycological Progress* 14:86.
- Yu Y, Cui Y, Hsiang T, Zeng ZQ, Yu ZH. 2015. Isolation and identification of endophytes from roots of *Cymbidium goeringii* and *Cymbidium faberi* (Orchidaceae). *Nova Hedwigia* 1001:57–64.
- Zerbino DR. 2010. Using the velvet de novo assembler for short-read sequencing technologies. *Current Protocols in Bioinformatics*. Chapter 11: Unit 11.5.

LEGENDS

FIG. 1. *Bifiguratus adelaidae* colony morphology on different media and temperatures. A. MEA+ at 25 °C. B. SDA+ at 25 °C. C. EYSA+ at 25 °C. D. PYG+ at 25 °C. E. MEA+ at 35 °C. F. SDA+ at 35 °C. G. EYSA+ at 35 °C. H. PYG+ at 35 °C. Bars= 10 mm. Images presented belong to

different isolates. All isolates (AZ0501, TLT265, YP243) showed consistent growth in each media.

FIG. 2. Radial growth rates of *B. adelaidae* per day on MEA+ under three different temperatures during 30 days. Vertical bars indicate standard deviation. A. Isolate TLT265. B. Isolate YP243. C. Isolate AZ0501, triangle = 10 °C circle = 25 °C box= 35 °C.

FIG. 3. Light microscopy of *B. adelaidae* isolate TLT265. A. Yeast cells. B. Putative intercalary chlamydospores (dark purple) in coenocytic hyphae. C. Sporangium. Bars= 25 µm.

FIG. 4. Scanning electron micographs of *B. adelaidae* hyphal (A–C) and yeast (D–F) structures.

A. Putative germinating chlamydospore with several germ tubes. B. Profuse vegetative hyphae C. Hyphal wall surface ornamentations. D. Yeast bud scar (arrow). E. Budding of yeast cells (arrow). F. Germinating yeast cells, arrows point to germ tubes. Micrograph E is from YP243; all other images are from isolate TLT265.

FIG. 5. Phylogenetic placement of *Bifiguratus adelaidae* within Mucoromycota (Spatafora et al. 2016) based upon concatenated 18S and 28S rRNA gene sequence alignment. *Basidobolus ranarum* was used as outgroup. Branches with Bayesian posterior probabilities values >0.95 are thickened and ML bootstrap support values >70 are shown.

FIG. 6. RAxML phylogenetic tree of Mucoromycota based on a concatenated alignment of 434 conserved orthologous proteins. *Bifiguratus adelaidae* isolate AZ0501 is in bold. Bootstrap values are shown on respective branches.

FIG. 7. Bacteria associated with isolates of *B. adelaidae*. A. Sequence read abundance of the 25 bacterial operational taxonomic units (OTUs) detected through the 454 pyrosequencing of three isolates of *B. adelaidae* grown at different temperatures and on different media types. OTU relative sequence abundance is expressed as log10 of read number. The plot was generated with

the R package “phyloseq” (McMurdie and Holmes 2013). Bars with less than one read in a single samples group are not displayed in bar plots. B. Venn diagrams of the bacterial OTUs comparing diversity between isolates, media, and temperature (left to right). C, D. Scanning electron micrographs of bacteria present on *B. adalaidae* isolate AZ0501. Bars= 5 µm.

TABLES

TABLE 1. Benchmarking universal single-copy orthologs (BUSCO) results for *Bifiguratus adalaidae* and other Mucoromycotina.

SUPPLEMENTARY MATERIAL

TABLE S1. Estimates of evolutionary divergence between sequences of ITS, 28S, and 18S nrRNA for isolates AZ0501, TLT265, and YP243 of *Bifiguratus adalaidae*.

TABLE S2. Statistics of the completeness of the *Bifiguratus adalaidae* genome using CEGMA based on 248 CEGs

TABLE S3. Isolation of bacteria associated with *Bifiguratus adalaidae*.

TABLE S4. Bacterial operational taxonomic units (OTUs) detected by 454 pyrosequencing on isolates of *Bifiguratus adalaidae* under different media and temperature treatments.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

